

Wright State University

CORE Scholar

---

Symposium of Student Research, Scholarship,  
and Creative Activities Materials

Office of the Vice Provost for Research

---

4-15-2016

## Investigating the After-Effects of Stochastic Resonance for Use in Laparoscopic Surgery

Muhammad S. Hamdan

Wright State University - Main Campus, hamdan.6@wright.edu

Caroline G. L. Cao

Wright State University - Main Campus, caroline.cao@wright.edu

Follow this and additional works at: [https://corescholar.libraries.wright.edu/urop\\_celebration](https://corescholar.libraries.wright.edu/urop_celebration)



Part of the [Arts and Humanities Commons](#), [Engineering Commons](#), [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), [Physical Sciences and Mathematics Commons](#), and the [Social and Behavioral Sciences Commons](#)

---

### Repository Citation

Hamdan , M. S., & Cao , C. G. (2016). *Investigating the After-Effects of Stochastic Resonance for Use in Laparoscopic Surgery*. .

This Abstract is brought to you for free and open access by the Office of the Vice Provost for Research at CORE Scholar. It has been accepted for inclusion in Symposium of Student Research, Scholarship, and Creative Activities Materials by an authorized administrator of CORE Scholar. For more information, please contact [library-corescholar@wright.edu](mailto:library-corescholar@wright.edu).

# *Investigating the After-Effects of Stochastic Resonance for Use in Laparoscopic Surgery*

*Muhammed S. Hamdan & Caroline GL Cao*

*Ergonomics in Remote Environments Laboratory, WSU*

Laparoscopic surgery is a modern surgical technique in which surgeons insert tools through small incisions in the abdomen to perform a surgical procedure. Surgeons typically have only limited 2D visual feedback based on a small camera that is inserted into the abdomen, and minimal haptic feedback that is transmitted through the rigid surgical tools. It is difficult for surgeons to detect subthreshold haptic feedback that surgical instruments transfer onto their hands. Research has shown that Stochastic Resonance (SR), which is the phenomenon of introducing white noise into a nonlinear system to enhance the detection of subthreshold stimuli, is capable of increasing the sensitivity of surgeons' hands and, therefore, improving their detection of embedded tumors during simulated laparoscopic surgery. There is evidence to suggest that an after-effect of SR exists long after the application of the white noise. However, not much work has been done to quantify the extent of the SR after-effect on users, and specifically, on surgeons. Our objective is to measure the extent of SR after-effect after a prolonged SR application in the form of a vibration signal. The results obtained from this research will allow us to design an effective enabling technology that applies SR in clinical settings, enhancing the sensitivity and performance of surgeons in laparoscopic surgery.